

# Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field

---

**Andrei O. Barvinsky**

*Theory Department, Lebedev Physics Institute, Leninsky Prospekt 53, Moscow 119991, Russia*

*E-mail:* `barvin@td.lpi.ru`

**Alexander Yu. Kamenshchik**

*Dipartimento di Fisica and INFN, via Irnerio 46, 40126 Bologna, Italy  
L. D. Landau Institute for Theoretical Physics, Moscow 119334, Russia*

*E-mail:* `kamenshchik@bo.infn.it`

**Claus Kiefer**

*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Strasse 77, 50937 Köln, Germany*

*E-mail:* `kiefer@thp.uni-koeln.de`

**Alexei A. Starobinsky**

*L. D. Landau Institute for Theoretical Physics, Moscow 119334, Russia  
RESCEU, Graduate School of Science, The University of Tokyo, Tokyo 113-0033, Japan*

*E-mail:* `alstar@landau.ac.ru`

**Christian Steinwachs**

*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Strasse 77, 50937 Köln, Germany*

*E-mail:* `cst@thp.uni-koeln.de`

**ABSTRACT:** We consider the renormalization group improvement in the theory of the Standard Model Higgs boson playing the role of an inflaton with a strong non-minimal coupling to gravity. It suggests the range of the Higgs mass  $135.6 \text{ GeV} \lesssim M_H \lesssim 184.5 \text{ GeV}$  compatible with the current CMB data (the lower WMAP bound on  $n_s$ ), which is close to the widely accepted range dictated by the electroweak vacuum stability and perturbation theory bounds. We find the phenomenon of asymptotic freedom induced by this non-minimal curvature coupling, which brings the theory to the weak coupling domain everywhere except at the lower and upper boundary of this range. The renormalization group running of the basic quantity  $\mathbf{A_I}$  – the anomalous scaling in the non-minimally coupled Standard Model, which analytically determines all characteristics of the CMB spectrum – brings  $\mathbf{A_I}$  to small negative values at the inflation scale. This property is crucial for the above results and may also underlie the formation of initial conditions for the inflationary dynamics in quantum cosmology.

**KEYWORDS:** Inflation, Higgs boson, Standard Model, Renormalization group.

---

## Contents

|                                                                                                           |           |
|-----------------------------------------------------------------------------------------------------------|-----------|
| <b>1. Introduction</b>                                                                                    | <b>1</b>  |
| <b>2. Anomalous scaling parameter in theories with a large non-minimal curvature coupling</b>             | <b>3</b>  |
| <b>3. One-loop approximation</b>                                                                          | <b>6</b>  |
| <b>4. RG improvement</b>                                                                                  | <b>8</b>  |
| <b>5. Inflationary stage versus post-inflationary running</b>                                             | <b>11</b> |
| <b>6. Numerical analysis</b>                                                                              | <b>14</b> |
| <b>7. Conclusions and discussion</b>                                                                      | <b>19</b> |
| <b>A. Renormalization of potential and non-minimal coupling terms – role of Higgs and Goldstone modes</b> | <b>24</b> |

---

## 1. Introduction

The task of constructing a fundamental particle model accounting for an inflationary scenario and its observationally consistent predictions for the CMB was undertaken in [1] and, more recently, in the series of papers [2, 3, 4, 5, 6]. While the relatively old work [1] had suggested that due to quantum effects inflation depends not only on the inflaton-graviton sector of the system but rather is strongly effected by its GUT contents, the paper by Bezrukov and Shaposhnikov [2] transcended this idea from the remote land of Grand Unification Theory to the seemingly firm ground of the Standard Model (SM) with the Higgs field playing the role of an inflaton. This has provoked new interest in a once rather popular [7, 8, 9, 10, 1, 11, 12] but then nearly forgotten model with the Lagrangian of the graviton-inflaton sector given by

$$L(g_{\mu\nu}, \Phi) = \frac{1}{2} (M_P^2 + \xi |\Phi|^2) R - \frac{1}{2} |\nabla \Phi|^2 - V(|\Phi|), \quad (1.1)$$

$$V(|\Phi|) = \frac{\lambda}{4} (|\Phi|^2 - v^2)^2, \quad |\Phi|^2 = \Phi^\dagger \Phi, \quad (1.2)$$

where  $\Phi$  is one of the scalar multiplets of the GUT-type sector of the system, whose expectation value plays the role of an inflaton and which has a strong non-minimal curvature coupling with  $\xi \gg 1$ . Here,  $M_P = m_P / \sqrt{8\pi} \approx 2.4 \times 10^{18}$  GeV is a reduced Planck mass,  $\lambda$  is a quartic self-coupling of  $\Phi$ , and  $v$  is a symmetry breaking scale.

The motivation for this model was based in the early papers [7, 8, 9, 10] on the observation that the problem of an exceedingly small quartic coupling  $\lambda \sim 10^{-13}$ , dictated by the amplitude of primordial scalar (density) perturbations [13, 14, 15], can be solved by using a non-minimally coupled inflaton with a large value of  $\xi$ , because the CMB anisotropy  $\Delta T/T \sim 10^{-5}$  is proportional to the ratio  $\sqrt{\lambda}/\xi$ , rather than to  $\sqrt{\lambda}$  as in the minimal case. Therefore, a small  $\Delta T/T$  can be obtained even for  $\lambda$  close to unity (but still small enough to justify perturbative expansion in  $\lambda$ ) if  $\xi \sim 10^4$ .

Later the model (1.1) with the GUT-type sector of matter fields was used to generate initial conditions for inflation [1] within the concept of the no-boundary [16] and tunneling cosmological states [17, 18, 19, 20]. The quantum evolution with these initial data was considered in [11, 21]. In particular, it was shown that quantum effects are critically important for this scenario.

A similar model but with the Standard Model (SM) Higgs boson  $\Phi$  playing the role of an inflaton instead of the abstract GUT setup of [1, 11] was suggested in [2]. This work advocated the consistency of the corresponding CMB data with WMAP observations. However, it disregarded quantum effects<sup>1</sup> which were later taken into account in [3] at the level of the one-loop approximation. Application of the general method of [1, 11] to the Standard Model Higgs inflation has led to the lower bound on the Higgs mass  $M_H \gtrsim 230$  GeV, originating from the observational restrictions on the CMB spectral index [3]. However, this pessimistic conclusion, which contradicts the widely accepted range  $115 \text{ GeV} \leq M_H \leq 180 \text{ GeV}$ , did not take into account  $O(1)$  contributions due to running logarithms in the quantum effective action, which can qualitatively improve the situation<sup>2</sup>. This was nearly simultaneously observed in [23, 4, 5] where the renormalization group (RG) improvement of the one-loop results of [3] was considered and it was found that the Higgs-mass range compatible with the CMB data almost coincides with the conventional one.

In contrast to the numerical and, therefore, not entirely transparent analysis of [4, 5], we develop here a method which retains the analytical formalism of [3] by incorporating the technique of running coupling constants. This supports the results of [3] and, moreover, explicitly reveals a mechanism which brings the CMB compatible range of the Higgs mass closer to the conventional Standard Model domain. This mechanism also explains the perfect efficiency of the perturbation theory at energy scales of the inflationary stage; since this is analogous to asymptotic freedom in QCD, we call it here asymptotic freedom, too. This mechanism is mediated by the effect of longitudinal (virtual) gravitons; gravitation thus serves as a regulator making the Standard Model sector softer.

The organization of the paper is as follows. In Sects. 2 and 3 we discuss the notion of anomalous scaling and recapitulate the results of the one-loop approximation of [3]. In Sect. 4 we develop the RG improvement for the effective action of the theory, present beta functions for running couplings with a special emphasis on loop contributions of Higgs

---

<sup>1</sup>This was caused by the use of a special field-dependent renormalization scheme [22] suppressing the contributions of running logarithms in the effective potential.

<sup>2</sup>The possibility of such contributions was, in fact, mentioned in the paper [3], but the sign of their effect was not predictable at the moment of publication.

particles and Goldstone modes in the presence of the non-minimal coupling to graviton [11, 3, 5]. In Sect. 5 we prove the applicability of the analytic method of [3] for the CMB dynamics at the inflationary stage, which is based on the shortness of this stage compared to the post-inflationary evolution to the present-day electroweak vacuum. Sect. 6 contains results of numerical simulations for the RG flow of coupling constants, the anomalous scaling  $\mathbf{A}(t)$  in particular, which impose CMB and Standard Model bounds on the Higgs mass and reveal the asymptotic-freedom mechanism based on the behavior of  $\mathbf{A}(t)$  and  $\lambda(t)$ . In concluding Sect. 7, we discuss the limitations of the obtained results associated with gauge and parametrization dependence of the underlying formalism and suggest the method of their resolution. We also compare in the context of these issues our conclusions with the results of [6] and [5]. We show that the RG treatment of the model is essentially more sensitive to the CMB bounds than in [6], especially at the upper limit of the Higgs mass range  $\simeq 185$  GeV. We also find that the RG improvement makes the scenario realistic in which the initial conditions for inflation are generated from quantum cosmology in the form of a sharp probability peak of the tunneling wavefunction [17, 18, 19, 20]. The Appendix contains the derivation of SM beta functions in the presence of a strong non-minimal curvature coupling of the Higgs field.

## 2. Anomalous scaling parameter in theories with a large non-minimal curvature coupling

The usual understanding of non-renormalizable theories is that renormalization of higher-dimensional operators does not effect the renormalizable sector of low-dimensional operators, because the former ones are suppressed by powers of a cutoff – the Planck mass  $M_P$  [24]. Therefore, beta functions of the Standard Model sector are not expected to be modified by graviton loops. The situation with the non-minimal coupling is more subtle. Due to the mixing of the Higgs scalar field with the longitudinal part of gravity in the kinetic term of the Lagrangian (1.1), an obvious suppression of pure graviton loops by the effective Planck mass,  $M_P^2 + \xi\varphi^2 \gg M_P^2$ , for large  $\xi$  proliferates to the sector of the Higgs field, so that certain parts of beta functions, which induce Landau poles, are strongly damped by large  $\xi$  [5]. Therefore, running coupling constants like  $\lambda(t)$  turn out to be small or remain finite at the inflation scale. In particular, a special combination of coupling constants  $\mathbf{A}$  which we call *anomalous scaling* [1] becomes very small and reduces the CMB-compatible Higgs-mass bound. The importance of  $\mathbf{A}$  follows from the fact observed in [1, 11, 3] that due to large  $\xi$ , quantum effects and their CMB manifestation are universally determined by  $\mathbf{A}$ . The nature of this quantity, which was initially introduced in the context of a generic gauge theory non-minimally coupled to gravity [1], is as follows.

Let the model contain in addition to (1.1) also a set of scalar fields  $\chi$ , vector gauge bosons  $A_\mu$  and spinors  $\psi$ , which have a typical interaction with  $\Phi$  dictated by the local gauge invariance. If we denote by  $\varphi$  the inflaton – the only nonzero component of the mean value of  $\Phi$  in the cosmological state,  $\varphi^2 = \Phi^\dagger\Phi$ , then the non-derivative part of the

interaction of these particles with  $\varphi$  can be schematically written down as

$$\mathbf{L}_{\text{int}} = - \sum_{\chi} \frac{1}{2} \lambda_{\chi} \chi^2 \varphi^2 - \sum_A \frac{1}{2} g_A^2 A_{\mu}^2 \varphi^2 - \sum_{\psi} y_{\psi} \varphi \bar{\psi} \psi, \quad (2.1)$$

where  $\lambda_{\chi}$ ,  $g_A$  and  $y_{\psi}$  are the relevant constants of quartic, gauge, and Yukawa couplings.

On the background of a slowly varying inflaton  $\varphi$ , these terms generate masses  $m(\varphi)$  for all these particles, which are proportional to  $\varphi$ ,  $m(\varphi) \sim \varphi$ , and which in turn generate at one-loop order the Coleman-Weinberg potential

$$\sum_{\text{particles}} (\pm 1) \frac{m^4(\varphi)}{64\pi^2} \ln \frac{m^2(\varphi)}{\mu^2} = \frac{\lambda \mathbf{A}}{128\pi^2} \varphi^4 \ln \frac{\varphi^2}{\mu^2} + \dots \quad (2.2)$$

Here, the summation over particles includes the statistics as well as the sum over polarizations, and the overall coefficient  $\mathbf{A}$  schematically reads

$$\mathbf{A} = \frac{2}{\lambda} \left( \sum_{\chi} \lambda_{\chi}^2 + 3 \sum_A g_A^4 - 4 \sum_{\psi} y_{\psi}^4 \right), \quad (2.3)$$

where the coefficients 1, 3 and 4 in front of the sums are the number of degrees of freedom of a real scalar field, a massive vector field, and a charged Dirac spinor<sup>3</sup>. This quantity is, of course, well known, because as a coefficient of the logarithm it comprises the ultraviolet renormalization of  $\lambda$  and constitutes the conformal anomaly, or anomalous scaling, associated with the normalization scale  $\mu$  in (2.2). It includes the one-loop contributions of all particles except the graviton and the inflaton field  $\varphi$  itself, because their contributions are strongly suppressed by inverse powers of  $\xi \gg 1$  [11, 3, 5] – a property which will be discussed in much detail below.

The role of (2.3) is two-fold. First, in the context of the no-boundary and tunneling initial conditions [16, 17, 18, 19, 20] extended to the one-loop level [1], it appears in the distribution function of the quasi-de Sitter cosmological instantons  $\rho(\varphi)$  describing a quantum distribution of cosmological models with different initial  $\varphi$ ,

$$\rho(\varphi) \sim \left( \frac{\varphi}{\mu} \right)^{-6\mathbf{A}\xi^2/\lambda}. \quad (2.4)$$

This scaling (which gave rise to the name “anomalous scaling”) makes this distribution normalizable at  $\varphi \rightarrow \infty$  for positive  $\mathbf{A}$  [25] and, moreover, for the case of the tunneling cosmological wavefunction it generates a sharp probability peak in  $\rho(\varphi)$  at the following value of the inflaton field: the quantum scale of inflation derived in [1],  $\varphi_I^2 = 64\pi^2 M_P^2 / \xi \mathbf{A}$ .

Secondly, the anomalous scaling for  $\xi \gg 1$  determines the quantum rolling force in the effective equation of the inflationary dynamics [11, 21] and, consequently, yields the parameters of the CMB generated during inflation [3]. These parameters explicitly depend

---

<sup>3</sup>For a generic model,  $(\lambda_{\chi}, g_A^2, y_{\psi})$  comprise coupling constant matrices which give rise to mass matrices and their traces in the actual expression for  $\mathbf{A}$ .

on  $\mathbf{A}$  and the e-folding number  $N$  of the first horizon crossing by the primordial cosmological perturbation of a given wavelength. This dependence for the spectral index  $n_s$ , which belongs to the interval  $0.94 < n_s(k_0) < 0.99$  (the combined WMAP+BAO+SN data at the pivot point  $k_0 = 0.002 \text{ Mpc}^{-1}$  corresponding to  $N \simeq 60$  [26, 27]), gives the range of the anomalous scaling  $-12 < \mathbf{A} < 14$  [3].

On the other hand, in the Standard Model  $\mathbf{A}$  is expressed in terms of the masses of the heaviest particles –  $W^\pm$  boson,  $Z$  boson and top quark,

$$m_W^2 = \frac{1}{4} g^2 \varphi^2, \quad m_Z^2 = \frac{1}{4} (g^2 + g'^2) \varphi^2, \quad m_t^2 = \frac{1}{2} y_t^2 \varphi^2, \quad (2.5)$$

and the mass of three Goldstone modes

$$m_G^2 = \frac{V'(\varphi)}{\varphi} = \lambda(\varphi^2 - v^2) \simeq \lambda\varphi^2. \quad (2.6)$$

Here,  $g$  and  $g'$  are the  $SU(2) \times U(1)$  gauge couplings,  $g_s$  is the  $SU(3)$  strong coupling and  $y_t$  is the Yukawa coupling for the top quark. We work in the transversal Landau gauge with no contribution to  $m_G^2$  from the gauge-fixing term. At the inflation stage with  $\varphi^2 \gg v^2$ , the Goldstone mass  $m_G^2$  is non-vanishing in contrast to its zero on-shell value in the electroweak vacuum [28]. It is again important to emphasize that the Higgs particle itself with the mass

$$m_H^2 = V''(\varphi) = \lambda(3\varphi^2 - v^2) \simeq 3\lambda\varphi^2 \quad (2.7)$$

does not contribute to  $\mathbf{A}$ , because its contribution is suppressed at the inflation scale by a small factor  $\sim 1/\xi^2$ ; this is shown in the Appendix.

Equation (2.2) then gives the expression

$$\mathbf{A} = \frac{3}{8\lambda} \left( 2g^4 + (g^2 + g'^2)^2 - 16y_t^4 \right) + 6\lambda. \quad (2.8)$$

In the conventional range of the Higgs mass  $115 \text{ GeV} \leq M_H \leq 180 \text{ GeV}$  [29], this quantity is at the electroweak scale in the range  $-48 < \mathbf{A} < -20$ , which strongly contradicts the CMB range given above and violates as well the normalizability of the distribution function (2.4). These two ranges can be brought together only by the price of raising  $\lambda \simeq M_H^2/2v^2$ , that is, by increasing the Higgs mass to 230 GeV [3].

Salvation comes, however, from the observation that the RG running of coupling constants is strong enough and drives  $\mathbf{A}$  to a range compatible with the CMB data *and* the conventional range of the Higgs mass. Thus, the formalism of [3] stays applicable but with the electroweak  $\mathbf{A}$  replaced by the running  $\mathbf{A}(t)$ ,

$$\mathbf{A}(t) = \frac{3}{8\lambda(t)} \left( 2g^4(t) + (g^2(t) + g'^2(t))^2 - 16y_t^4(t) \right) + 6\lambda(t). \quad (2.9)$$

Here,  $t = \ln(\varphi/\mu)$  is the running scale of the RG improvement of the effective potential [30], where  $\mu$  is a normalization point which we choose to coincide with the top quark mass  $\mu = M_t$  (we denote physical (pole) masses by capital letters in contrast to the running masses (2.5) above). In the following we shall present the details of this mechanism.

### 3. One-loop approximation

The inflationary model with a non-minimally coupled Higgs-inflaton was considered in the one-loop approximation in [3]. The low-derivative part of its effective action (appropriate for the inflationary slow-roll scenario),

$$S[g_{\mu\nu}, \varphi] = \int d^4x g^{1/2} \left( -V(\varphi) + U(\varphi) R(g_{\mu\nu}) - \frac{1}{2} G(\varphi) (\nabla\varphi)^2 \right), \quad (3.1)$$

contains coefficient functions which in the one-loop approximation read as

$$V(\varphi) = \frac{\lambda}{4}(\varphi^2 - \nu^2)^2 + \frac{\lambda\varphi^4}{128\pi^2} \mathbf{A} \ln \frac{\varphi^2}{\mu^2}, \quad (3.2)$$

$$U(\varphi) = \frac{1}{2}(M_P^2 + \xi\varphi^2) + \frac{\varphi^2}{32\pi^2} \left( \mathbf{C} \ln \frac{\varphi^2}{\mu^2} + D \right), \quad (3.3)$$

$$G(\varphi) = 1 + \frac{1}{32\pi^2} \left( F \ln \frac{\varphi^2}{\mu^2} + E \right). \quad (3.4)$$

These functions contain numerical coefficients  $\mathbf{A}, \mathbf{C}, D, F, E$  determined by contributions of quantum loops of all particles and include the dependence on the UV normalization scale  $\mu$ . For  $\xi \gg 1$  the inflationary stage and the corresponding CMB parameters critically depend only on the anomalous scaling  $\mathbf{A}$  defined by (2.2) [1, 11, 21, 3] and the part of  $\mathbf{C}$  which is linear in  $\xi$ . As shown in Appendix A, the latter is contributed by the Goldstone mass  $m_G^2$  and equals

$$\mathbf{C} = 3\xi\lambda + O(\xi^0). \quad (3.5)$$

In fact, this expression yields a quantum correction to the non-minimal coupling  $\xi$ , as can be seen from (3.3). Other coefficients, the normalization scale  $\mu$  inclusive, are irrelevant in the leading order of the slow roll expansion.

Inflation and its CMB are easy to analyze in the Einstein frame of fields  $\hat{g}_{\mu\nu}$ ,  $\hat{\varphi}$ , which is related to the Jordan frame used in (3.1) by the equations

$$\hat{g}_{\mu\nu} = \frac{2U(\varphi)}{M_P^2} g_{\mu\nu}, \quad \left( \frac{d\hat{\varphi}}{d\varphi} \right)^2 = \frac{M_P^2}{2} \frac{GU + 3U'^2}{U^2}. \quad (3.6)$$

The action (3.1) in the Einstein frame,  $\hat{S}[\hat{g}_{\mu\nu}, \hat{\varphi}] = S[g_{\mu\nu}, \varphi]$ , has a minimal coupling,  $\hat{U} = M_P^2/2$ , a canonically normalized inflaton field,  $\hat{G} = 1$ , and a new inflaton potential,

$$\hat{V}(\hat{\varphi}) = \left( \frac{M_P^2}{2} \right)^2 \frac{V(\varphi)}{U^2(\varphi)} \Big|_{\varphi=\varphi(\hat{\varphi})}. \quad (3.7)$$

In view of (3.2), (3.3) and (3.5), at the inflation scale with  $\varphi > M_P/\sqrt{\xi} \gg v$  and  $\xi \gg 1$ , this potential reads

$$\hat{V} = \frac{\lambda M_P^4}{4\xi^2} \left( 1 - \frac{2M_P^2}{\xi\varphi^2} + \frac{\mathbf{A}_I}{16\pi^2} \ln \frac{\varphi}{\mu} \right), \quad (3.8)$$



where the parameter  $\mathbf{A_I}$  represents the anomalous scaling (2.8) modified by quantum corrections to  $U$  – the renormalization of the non-minimal term (3.5),

$$\mathbf{A_I} = \mathbf{A} - 12\lambda = \frac{3}{8\lambda} \left( 2g^4 + (g^2 + g'^2)^2 - 16y_t^4 \right) - 6\lambda. \quad (3.9)$$

This quantity – which we will call *inflationary anomalous scaling* – enters the expressions for the standard slow-roll parameters,

$$\hat{\varepsilon} \equiv \frac{M_P^2}{2} \left( \frac{1}{\hat{V}} \frac{d\hat{V}}{d\hat{\varphi}} \right)^2 = \frac{4}{3} \left( \frac{M_P^2}{\xi \varphi^2} + \frac{\mathbf{A_I}}{64\pi^2} \right)^2, \quad (3.10)$$

$$\hat{\eta} \equiv \frac{M_P^2}{\hat{V}} \frac{d^2 \hat{V}}{d\hat{\varphi}^2} = -\frac{4M_P^2}{3\xi \varphi^2}, \quad (3.11)$$

and ultimately determines all the characteristics of inflation. In particular, the smallness of  $\hat{\varepsilon}$  yields the range of the inflationary stage  $\varphi > \varphi_{\text{end}}$ , terminating at a value of  $\hat{\varepsilon}$  which for elegance of the formalism we chose to be  $\hat{\varepsilon}_{\text{end}} = 3/4$ . Then the inflaton value at the exit from inflation equals  $\varphi_{\text{end}} \simeq 2M_P/\sqrt{3\xi}$  under the natural assumption that the perturbation loop expansion is applicable for  $\mathbf{A_I}/64\pi^2 \ll 1$ . The duration of inflation which starts at  $\varphi$  takes in units of the scale factor e-folding number  $N$  a particularly simple form [3],

$$\frac{\varphi^2}{\varphi_I^2} = e^x - 1, \quad (3.12)$$

$$\varphi_I^2 = \frac{64\pi^2 M_P^2}{\xi \mathbf{A_I}}, \quad (3.13)$$

where the quantum scale of inflation  $\varphi_I$  and a special parameter  $x$  containing the e-folding number,

$$x \equiv \frac{N \mathbf{A_I}}{48\pi^2}, \quad (3.14)$$

directly involve the anomalous scaling  $\mathbf{A_I}$  which is essentially a quantum quantity.

This relation determines the Fourier power spectrum for the scalar metric perturbation  $\zeta$ ,  $\Delta_\zeta^2(k) \equiv \langle k^3 \zeta_{\mathbf{k}}^2 \rangle = \hat{V}/24\pi^2 M_P^4 \hat{\varepsilon}$ , where the right-hand side is taken at the first horizon crossing,  $k = aH$ , relating the comoving perturbation wavelength  $k^{-1}$  to the e-folding number  $N$ ,

$$\Delta_\zeta^2 = \frac{N^2}{72\pi^2} \frac{\lambda}{\xi^2} \left( \frac{e^x - 1}{x e^x} \right)^2. \quad (3.15)$$

The CMB spectral index  $n_s \equiv 1 + d \ln \Delta_\zeta^2 / d \ln k = 1 - 6\hat{\varepsilon} + 2\hat{\eta}$ , the tensor to scalar ratio  $r = 16\hat{\varepsilon}$  and the spectral index running  $\alpha \equiv dn_s/d \ln k \simeq -dn_s/dN$  correspondingly read as

$$n_s = 1 - \frac{2}{N} \frac{x}{e^x - 1}, \quad (3.16)$$

$$r = \frac{12}{N^2} \left( \frac{x e^x}{e^x - 1} \right)^2, \quad (3.17)$$

$$\alpha = -\frac{2}{N^2} \frac{x^2 e^x}{(e^x - 1)^2}. \quad (3.18)$$

Note that for  $|x| \ll 1$  these predictions exactly coincide with those [32, 33] of the  $f(R) = M_P^2(R + R^2/6M^2)/2$  inflationary model [34] with the scalar particle (scalaron) mass  $M = M_P\sqrt{\lambda}/\sqrt{3}\xi$ .

With the spectral index constraint  $0.94 < n_s(k_0) < 0.99$  at the pivot point  $k_0 = 0.002 \text{ Mpc}^{-1}$  corresponding to  $N \simeq 60$  [26, 27] these relations immediately give the range of anomalous scaling  $-12 < \mathbf{A}_I < 14$ . As mentioned above, this contradicts the Standard Model bound  $-48 < \mathbf{A}_I < -20$  existing in the conventional range of the Higgs mass  $115 \text{ GeV} \leq M_H \leq 180 \text{ GeV}$  [29].<sup>4</sup>

The limitation of this approximation follows from the fact that, despite the smallness of  $\mathbf{A}_I/64\pi^2$ , the running of the logarithm from the electroweak scale  $\varphi = v$  (where the Higgs and other particle masses are determined) to the inflation scale  $\varphi \simeq |\varphi_I|$  is big enough and changes the predictions by  $O(1)$  factors. Indeed, with the values of  $\xi \sim 10^4$  and  $|\mathbf{A}_I| \sim 10$  dictated by the results of [3], the estimate for the logarithmic running  $\ln(|\varphi_I|^2/v^2) \sim 60$  yields the contribution

$$\frac{\mathbf{A}_I}{64\pi^2} \ln \frac{|\varphi_I|^2}{v^2} \sim 2 = O(1), \quad (3.19)$$

which requires resummation within the RG improvement. The necessity of this improvement was mentioned in [3], and it was realized in [4, 5]. Below we develop a combined numerical and analytical scheme of this improvement, which recovers the above formalism of Eqs. (3.12)–(3.18) incorporating running coupling constants.

#### 4. RG improvement

According to the Coleman–Weinberg technique [30], the one-loop RG improved effective action has the form (3.1) with the coefficients

$$V(\varphi) = \frac{\lambda(t)}{4} Z^4(t) \varphi^4, \quad (4.1)$$

$$U(\varphi) = \frac{1}{2} \left( M_P^2 + \xi(t) Z^2(t) \varphi^2 \right), \quad (4.2)$$

$$G(\varphi) = Z^2(t). \quad (4.3)$$

Here,  $t = \ln(\varphi/M_t)$  is the running RG scale<sup>5</sup>, and the running couplings  $\lambda(t)$ ,  $\xi(t)$  and the field renormalization  $Z(t)$  incorporate a summation of powers of logarithms and belong to the solution of the RG equations

$$\frac{dg_i}{dt} = \beta_{g_i}, \quad g_i = (\lambda, \xi, g, g', g_s, y_t), \quad (4.4)$$

$$\frac{dZ}{dt} = \gamma Z \quad (4.5)$$

<sup>4</sup>These estimates were obtained in [3] for the parameter  $\mathbf{A}$  disregarding the contribution of Goldstone modes ( $+6\lambda$  and  $-6\lambda$  terms in (2.8) and (3.9), respectively), but numerically with  $\lambda \lesssim 1$  the inclusion of this contribution leads to qualitatively the same conclusions.

<sup>5</sup>Application of the Coleman–Weinberg technique removes the ambiguity in the choice of the RG scale in cosmology – an issue discussed in [31].

for the full set of coupling constants in the “heavy” sector of the model with the corresponding beta functions  $\beta_{g_i}$  and the anomalous dimension  $\gamma$  of the Higgs field.<sup>6</sup> This sector of the Standard Model includes the  $SU(2) \times U(1)$  gauge couplings  $g$  and  $g'$ , the  $SU(3)$  strong coupling  $g_s$ , and the Yukawa coupling  $y_t$  for the top quark.

An important subtlety with these  $\beta$  functions is the effect of the non-minimal curvature coupling of the Higgs field. For large  $\xi$  the kinetic term of the tree-level action has a strong mixing between the graviton  $h_{\mu\nu}$  and the quantum part of the Higgs field  $\sigma$  on the background  $\varphi$ . Symbolically it has the structure

$$(M_P^2 + \xi^2 \varphi^2) h \nabla \nabla h + \xi \varphi \sigma \nabla \nabla h + \sigma \square \sigma,$$

which yields a propagator whose elements are suppressed by a small  $1/\xi$ -factor in all blocks of the  $2 \times 2$  graviton-Higgs sector. For large  $\varphi \gg M_P/\sqrt{\xi}$ , the suppression of pure graviton loops is, of course, obvious because the effective Planck-mass squared exceeds by far the standard Planck-mass squared,  $M_P^2 + \xi \varphi^2 \gg M_P^2$ . Due to the mixing, this suppression proliferates to the full graviton-Higgs sector of the theory. To make this statement quantitative, one can go over to the Einstein frame (3.6) in which the propagator is diagonal and canonically normalized in the space of perturbations of  $\hat{g}_{\mu\nu}$  and  $\hat{\varphi}$  (in the background covariant DeWitt gauge). The above propagator can be uplifted back to the Einstein frame by matrix multiplication with the Jacobian matrices  $\partial(g_{\mu\nu}, \varphi)/\partial(\hat{g}_{\alpha\beta}, \hat{\varphi})$ . Its Higgs-Higgs block, in particular, gets suppressed by the factor  $s(\varphi)$

$$\left(\frac{\partial \varphi}{\partial \hat{\varphi}}\right)^2 \frac{1}{\hat{g}^{1/2}(\hat{\square} - \hat{m}_H^2)} = \frac{s(\varphi)}{g^{1/2}(\square - m_H^2)}, \quad (4.6)$$

$$s(\varphi) \equiv \frac{U}{GU + 3U'^2} = \frac{M_P^2 + \xi \varphi^2}{M_P^2 + (6\xi + 1)\xi \varphi^2}, \quad (4.7)$$

where we took into account the scaling  $\hat{g}^{1/2}(\hat{\square} - \hat{m}_H^2) = (2U/M_P^2)g^{1/2}(\square - m_H^2)$  under the conformal transformation (3.6) (disregarding spacetime gradients of  $\varphi$  and bearing in mind conformal rescaling of particle masses [3]). The origin of this suppression factor  $s(\varphi)$  within the original Jordan frame is clearly shown in Appendix A.

This mechanism was first understood in [9] when estimating the CMB generation in a non-minimal inflationary model. It was considered in [35] in the context of the generalized RG approach to nonrenormalizable theories. Also it justifies the omission of graviton loops [11, 21] and modifies the beta functions of the SM sector of this theory [5] at the high energy scale relevant for inflation. This modification is due to the suppression of every Higgs propagator (4.6) by the factor  $s(\varphi)$  [5], which is very small for  $\varphi \gg M_P/\sqrt{\xi}$ ,  $s \simeq 1/6\xi$ , but tends to one in the vicinity of the electroweak scale  $v \ll M_P/\xi$ . Such a

---

<sup>6</sup>Note that running couplings in the effective potential theory are the characteristics of the RG equation in partial derivatives for this potential, rather than the usual running coupling constants directly run by  $\beta$ 's. Thus the right-hand sides of RG eqs.(4.4)-(4.5) should contain the factor  $1/(1 - \gamma)$  [30] which we discard in the one-loop RG improvement.

modification justifies, in fact, the extension beyond the scale  $M_P/\xi$  interpreted in [36, 37] as a natural validity cutoff of the theory<sup>7</sup>.

Of course, the factor  $s(\varphi)$  makes the counterterms of the theory nonpolynomial in  $\varphi$ , and the theory as a whole becomes nonrenormalizable. One might think that this completely invalidates the construction of the usual RG improvement and requires the generalization of the latter to RG with an infinite set of charges (perhaps of the functional nature like in [39, 35]). However, there exists a shortcut to a simpler formulation based on the structure of the factor (4.7). In the logarithmic scale of the variable  $t = \ln(\varphi/\mu)$ , strongly compressing the transition domain  $M_P/\xi \lesssim \varphi \lesssim M_P/\sqrt{\xi}$  for  $s(\varphi)$  from 1 to  $1/6\xi \simeq 0$ , this factor looks very much like a step function  $\theta(t_0 - t)$ ,  $t_0 \simeq \ln(\varphi_0/\mu) = \ln(M_P/\sqrt{6\xi}\mu)$ . So it is nearly constant for  $\varphi < \varphi_0$  and  $\varphi > \varphi_0$ . Therefore, in both phases of such an approximation the theory is renormalizable but has different running of coupling constants.<sup>8</sup> This qualitatively justifies the use of beta functions modified by  $s$ -factors smoothly interpolating between these two phases. Below we will see that numerical results for smooth and step function  $s$ -factors nearly coincide with one another.

There is an important subtlety with the modification of beta functions, which was disregarded in [5] (and the first version of this paper). The mnemonic rule of associating the factor  $s(\varphi)$  with every propagator of the Higgs multiplet, as suggested in [5], turned out to be incorrect. Goldstone modes, in contrast to the Higgs particle, are not coupled to curvature, and they do not have a kinetic term mixing with gravitons [6]. Therefore, their contribution is not suppressed by the  $s$ -factor of the above type. Separation of Goldstone contributions from the Higgs contribution leads to the following modification of the one-loop beta functions, which is essentially different from that of [5] (cf. also [40]):

$$\beta_\lambda = \frac{\lambda}{16\pi^2} (18s^2\lambda + \mathbf{A}(t)) - 4\gamma\lambda, \quad (4.8)$$

$$\beta_\xi = \frac{6\xi}{16\pi^2} (1 + s^2)\lambda - 2\gamma\xi, \quad (4.9)$$

$$\beta_{y_t} = \frac{y_t}{16\pi^2} \left( -\frac{2}{3}g'^2 - 8g_s^2 + \left(1 + \frac{s}{2}\right) y_t^2 \right) - \gamma y_t, \quad (4.10)$$

---

<sup>7</sup>The notion of a cutoff depends on the type of perturbation theory used. Typically it is determined for the case when all dimensional quantities – fields and their derivatives – are considered on equal footing and treated perturbatively. However, a nonperturbative summation of powers of the field for low-derivative processes (achieved by the conformal transformation to the Einstein frame) allows one to go beyond such a cutoff, and this is exactly the situation with the scales above  $M_P/\xi$ , see e.g. [38].

<sup>8</sup>This is analogous to the approach of [6] where the model was approximated by matching the usual low-energy SM phase with the chiral phase of the SM at the inflation energy scale. This picture, however, takes place in the Einstein frame of the theory and is aggravated by the problem of transition between two different parameterizations of one quantum theory — the Cartesian coordinates in the space of the Higgs multiplet in the low-energy phase versus spherical coordinates in the chiral phase of the SM. This transition is physically nontrivial and is discussed below in more detail.

and

$$\beta_g = -\frac{39-s}{12} \frac{g^3}{16\pi^2}, \quad (4.11)$$

$$\beta_{g'} = \frac{81+s}{12} \frac{g'^3}{16\pi^2}, \quad (4.12)$$

$$\beta_{g_s} = -\frac{7g_s^3}{16\pi^2}. \quad (4.13)$$

Here, the anomalous dimension of the Higgs field  $\gamma$  is given by a standard expression in the Landau gauge,

$$\gamma = \frac{1}{16\pi^2} \left( \frac{9g^2}{4} + \frac{3g'^2}{4} - 3y_t^2 \right), \quad (4.14)$$

the anomalous scaling  $\mathbf{A}(t)$  is defined by (2.9), and we have retained only the leading terms in  $\xi \gg 1$ . It will be important in what follows that this anomalous scaling contains the Goldstone contribution  $6\lambda$ , so that the full  $\beta_\lambda$  in (4.8) has a  $\lambda^2$ -term unsuppressed by  $s(\varphi)$  at large scale  $t = \ln(\varphi/\mu)$ . In Appendix A we derive the Goldstone and Higgs contributions to  $\beta_\lambda$  and  $\beta_\xi$ , which demonstrates their different suppression mechanisms under non-minimal coupling with curvature.

## 5. Inflationary stage versus post-inflationary running

As we will see, the inflationary stage in units of Higgs-field e-foldings is very short. This allows us to make one more approximation: we shall consider the solutions of the RG equations only up to terms linear in  $\Delta t \equiv t - t_{\text{end}} = \ln(\varphi/\varphi_{\text{end}})$ , where we choose as the initial data point the end of inflation  $t_{\text{end}}$ . This approximation will be justified later in most of the Higgs-mass range compatible with the CMB data.

Thus, we use beta functions (4.8) and (4.9) with  $s = 0$  to obtain

$$\lambda(t) = \lambda_{\text{end}} \left( 1 - 4\gamma_{\text{end}}\Delta t + \frac{\mathbf{A}(t_{\text{end}})}{16\pi^2} \Delta t \right), \quad (5.1)$$

$$\xi(t) = \xi_{\text{end}} \left( 1 - 2\gamma_{\text{end}}\Delta t + \frac{6\lambda}{16\pi^2} \Delta t \right). \quad (5.2)$$

Here,  $\lambda_{\text{end}}$ ,  $\gamma_{\text{end}}$ ,  $\xi_{\text{end}}$  are determined at  $t_{\text{end}}$  and  $\mathbf{A}_{\text{end}} = \mathbf{A}(t_{\text{end}})$  is also the particular value of the running anomalous scaling (2.9) at the end of inflation.

On the other hand, the RG improvement of the effective action (4.1)–(4.3) implies that this action coincides with the tree-level action with running couplings as functions of  $t = \ln(\varphi/\mu)$  for a new field

$$\phi = Z(t) \varphi. \quad (5.3)$$

This is because the running of  $Z(t)$  is slow,  $\partial_\mu \phi = Z(1 + \gamma)\partial_\mu \varphi \simeq Z\partial_\mu \varphi$ , and the kinetic term of the effective action in terms of  $\phi$  gets canonically normalized. Then, in view of (4.1)–(4.2) the RG improved potential for this field takes at the inflation stage the form

$$\hat{V} = \left( \frac{M_P^2}{2} \right)^2 \frac{V}{U^2} \simeq M_P^4 \frac{\lambda_{\text{end}}}{4\xi_{\text{end}}^2} \left( 1 - \frac{2M_P^2}{\xi_{\text{end}}\phi^2} + \frac{\mathbf{A}_I(t_{\text{end}})}{16\pi^2} \ln \frac{\phi}{\phi_{\text{end}}} \right), \quad (5.4)$$

where for the same reason we disregarded the running of  $Z$  in  $\ln(Z/Z_{\text{end}})$ . This is nothing but the one-loop potential (3.8) for the field  $\phi$  with a particular choice of the normalization point  $\mu = \phi_{\text{end}}$  and the couplings replaced by the values of the running ones at  $t_{\text{end}}$ .

This means that the formalism of [3] can directly be applied to determine the parameters of the CMB. They are mainly determined by the anomalous scaling  $\mathbf{A}_I$ , but now this quantity should be taken at  $t_{\text{end}}$  rather than at the electroweak scale  $t = 0$ . This can solve the problem of matching the electroweak range of the anomalous scaling  $-48 < \mathbf{A}_I(0) < -20$  (corresponding to  $115 \text{ GeV} \leq M_H \leq 180 \text{ GeV}$ ) with the range required by the CMB data,  $-12.4 < \mathbf{A}_I(t_{\text{end}}) < 14.1$  [3], because  $\mathbf{A}_I(t_{\text{end}}) \neq \mathbf{A}_I(0)$  due to the running.

To find  $\mathbf{A}_I(t_{\text{end}})$  and other couplings at  $t_{\text{end}}$  we have to obtain the RG flow interpolating between the  $t = 0$  and  $t_{\text{end}} \simeq 33$ . In contrast to the inflationary stage, the post-inflationary running is very large, because of its long duration, and requires numerical simulation. In order to specify more precisely the initial conditions at the top-quark scale,  $t = 0$ , we take into account the pole mass matching scheme relating the observable physical masses  $M_H$  and  $M_t$  to the running masses  $m_H = \sqrt{2\lambda(t)}v$  and (2.5) (and relevant couplings) [41, 42, 43]. One might think that these subtleties essentially exceed the precision of possible SM implications in early cosmology. However, the instability bound on the Higgs mass and its lower bound from the CMB data, which turn out to be very close to one another, are very sensitive to the pole mass matching effect, so we include them in numerical simulations.

To have a better comparison with [5, 6], we fix the  $t = 0$  initial conditions for the RG equation (4.4)-(4.5) at the top-quark scale  $M_t = 171 \text{ GeV}$ . For the weak interaction constants  $g, g'$  and the strong interaction constant  $g_s$ , they read [29]

$$g^2(0) = 0.4202, \quad g'^2(0) = 0.1291, \quad g_s^2(0) = 1.3460, \quad (5.5)$$

where  $g^2(0)$  and  $g'^2(0)$  are obtained by a simple one-loop RG flow from the conventional values of  $\alpha(M_Z) \equiv g^2/4\pi = 0.0338$ ,  $\alpha'(M_Z) \equiv g'^2/4\pi = 0.0102$  at the  $M_Z$ -scale, and the value  $g_s^2(0)$  at  $M_t$  is generated by the numerical program of [44]<sup>9</sup>. For the Higgs self-interaction constant  $\lambda$  and for the Yukawa top quark interaction constant  $y_t$  the initial conditions are determined by the pole mass matching scheme originally developed in [41, 42] and presented in the following form in the Appendix of [43]:

$$\lambda(0) = \frac{M_H^2}{2v^2}(1 + 2\Delta_H(M_H)), \quad (5.6)$$

$$y_t(0) = \frac{\sqrt{2}M_t}{v}(1 + \Delta_t(M_H)). \quad (5.7)$$

Here, the electroweak vacuum expectation value for the Higgs field is  $v = 246.22 \text{ GeV}$ , and  $\Delta_H(M_H)$  and  $\Delta_t(M_H)$  comprise relevant mass operator corrections in the effective Higgs and top quark propagators (cf. the relation (2.5) for masses without these corrections), which together with  $M_H$  depend on  $M_t$ ,  $M_Z$  and the Weinberg angle.

---

<sup>9</sup>The analytical algorithm of transition between different scales for  $g_s^2$  was presented in [45].

The initial condition  $\xi(0)$  is not directly known. It should be determined from the CMB normalization condition for the amplitude of the power spectrum (3.15)  $\Delta_\zeta^2 \simeq 2.5 \times 10^{-9}$  at the pivot point  $k_0 = 0.002 \text{ Mpc}^{-1}$  [26, 27] which we choose to correspond to  $N \simeq 60$ . As quantum perturbations of the observable  $\varphi$  and the new field (5.3) are obviously related by  $\zeta_\varphi = \zeta_\phi/Z$ , Eq. (3.15) immediately yields the following estimate on the ratio of coupling constants:

$$\frac{1}{Z_{\text{in}}^2} \frac{\lambda_{\text{in}}}{\xi_{\text{in}}^2} \simeq 0.5 \times 10^{-9} \left( \frac{x_{\text{in}} \exp x_{\text{in}}}{\exp x_{\text{in}} - 1} \right)^2 \quad (5.8)$$

at the moment of the first horizon crossing for  $N = 60$ , which we call the “beginning” of inflation and label by  $t_{\text{in}}$ .<sup>10</sup> This moment, in turn, can be determined from the equations (3.12)–(3.14) and (3.13) relating the value of the inflaton field at this moment  $\varphi_{\text{in}}$  to the e-folding number  $N$  and anomalous scaling  $\mathbf{A}_I(t_{\text{in}})$ . With  $\varphi_{\text{in}} = M_t \exp(x_{\text{in}})$ , this relation takes the form

$$t_{\text{in}} = \ln \frac{M_P}{M_t} + \frac{1}{2} \ln \frac{4N}{3\xi_{\text{in}}} + \frac{1}{2} \ln \frac{\exp x_{\text{in}} - 1}{x_{\text{in}}}. \quad (5.9)$$

The RG equations (4.4) for the six couplings  $(g, g', g_s, y_t, \lambda, \xi)$  with five initial conditions (5.5)–(5.7) and the final condition (5.8) at  $t_{\text{in}}$  defined by Eq.(5.9) uniquely determine the RG flow for given values of the Higgs and top quark masses. Boundary conditions for this flow are rather involved, because they go beyond a usual Cauchy problem and even the definition of the final moment (5.9) includes the whole history from  $t = 0$  to  $t = t_{\text{in}}$ , that is, they are nonlocal in  $t$ . By iterations, however, this problem can be numerically solved with *Mathematica*; essential simplifications originate from the fact that  $x_{\text{in}}$  turns out to be  $O(1)$  (more precisely  $\simeq 2$ , see below) and that the last term in (5.9) almost always can be discarded from the expression for  $t_{\text{in}}$  on top of the contribution  $\ln(M_P/M_t) \simeq 37$ .

The RG flow covers also the inflationary stage from the chronological end of inflation  $t_{\text{end}}$  to  $t_{\text{in}}$  (remember that the arrows of the RG parameter  $t$  and the physical time are opposite). As was mentioned above, at the end of inflation we choose the value  $\hat{\varepsilon} = 3/4$  of the slow roll parameter (3.10), and  $\varphi_{\text{end}} = M_P \sqrt{4/3\xi_{\text{end}}}$  under the assumption  $\mathbf{A}_I(t_{\text{end}})/64\pi^2 \ll 1$  which will be justified below. Therefore,

$$t_{\text{end}} = \ln \frac{M_P}{M_t} + \frac{1}{2} \ln \frac{4}{3\xi_{\text{end}}}. \quad (5.10)$$

Thus the duration of inflation in units of inflaton field e-foldings  $t_{\text{in}} - t_{\text{end}} = \ln(\varphi_{\text{in}}/\varphi_{\text{end}})$  is very short relative to the post-inflationary evolution  $t_{\text{end}} \sim 35$ ,

$$t_{\text{in}} - t_{\text{end}} = \frac{1}{2} \ln N + \frac{1}{2} \ln \frac{\xi_{\text{in}}}{\xi_{\text{end}}} + \frac{1}{2} \ln \frac{\exp x_{\text{in}} - 1}{x_{\text{in}}} \simeq \frac{1}{2} \ln N \sim 2, \quad (5.11)$$

where we took into account that  $\xi_{\text{in}} \simeq \xi_{\text{end}}$  and  $x = O(1)$ . This is, of course, typical for large-field inflationary models with  $N \sim \varphi^2/M_P^2$ .

This estimate gives the range of our approximation linear in logarithms. The linearization, say in (5.1), implies the bound  $|\mathbf{A}_I(t_{\text{end}})|\Delta t/16\pi^2 \ll 1$ , where  $\Delta t < t_{\text{in}} - t_{\text{end}} \simeq \ln N/2$ ,


<sup>10</sup>No modification of the spectral index due to  $Z$  occurs in the one-loop RG running, because the difference  $n_s^\varphi - n_s^\phi = 2\gamma dt/dN = (\gamma \mathbf{A}_I/48\pi^2)e^x/(e^x - 1)$  belongs to the two-loop order.

and this approximation holds for

$$\frac{|A_I(t_{\text{end}})|}{16\pi^2} \ll \frac{2}{\ln N} \simeq 0.5. \quad (5.12)$$

## 6. Numerical analysis

The running of  $A(t)$  strongly depends on the behavior of  $\lambda(t)$ . It is well known that for small Higgs masses the usual RG flow in SM leads to an instability of the electroweak vacuum caused by negative values of  $\lambda(t)$  in a certain range of  $t$ , see e.g. [46, 43] and references therein. The same happens with the modified RG for the non-minimally coupled Higgs field considered here.



**Figure 1:** Running  $\lambda(t)$  for five values of the Higgs mass above the instability threshold. Dashed curves mark the boundaries of the inflation domain  $t_{\text{end}} \leq t \leq t_{\text{in}}$ .

The numerical solution for  $\lambda(t)$  in the setting of the previous section is shown in Fig.1 for five values of the Higgs mass and the value of top-quark mass  $M_t = 171$  GeV. The lowest one corresponds to the critical value


$$M_H^{\text{inst}} \simeq 134.27 \text{ GeV}. \quad (6.1)$$

This is the boundary of the instability window for which  $\lambda(t)$  bounces back to positive values after vanishing at  $t_{\text{inst}} \sim 41.6$  or  $\varphi_{\text{inst}} \sim 80M_P$ .<sup>11</sup> It turns out that the corresponding  $\xi(t)$  is nearly constant and is about 5000 (see below), so that the factor (4.7) at  $t_{\text{inst}}$  is very small,  $s \simeq 1/6\xi \sim 0.00005$ . Thus the situation is different from the usual Standard Model with  $s = 1$ , and numerically the critical value turns out to be higher than the known SM stability bound  $\sim 125$  GeV [43].

<sup>11</sup>An over-Planckian scale of  $\varphi$  does not signify a breakdown of the semiclassical expansion, because the energy density  $\sim 10^{-10}M_P^4$  stays much below the Planckian value, see below.



Fig.1 shows that near the instability threshold  $M_H = M_H^{\text{inst}}$  the running coupling  $\lambda(t)$  stays very small for all scales  $t$  relevant to the observable CMB, (5.9)-(5.10). This follows from the fact that the positive running of  $\lambda(t)$  caused by the term  $(18s^2 + 6)\lambda^2$  in  $\beta_\lambda$ , (4.8), is much slower for  $s \ll 1$  than that of the usual SM driven by the term  $24\lambda^2$ . For larger  $M_H$  this suppression in the term  $18s^2\lambda^2$  is responsible for the origin of the bump on the plots at  $t \sim 26$  where the value of  $s$  drops from 1 to 0 and  $\lambda(t)$  continues growing but with a slower rate.




**Figure 2:** Running anomalous scaling for the critical Higgs mass (the red curve with a vertical segment at the singularity with  $t_{\text{inst}} \sim 41.6$ ) and for two masses in the stability domain (blue and green curves).

The smallness of  $\lambda$  improves, of course, the efficiency of the perturbation theory. On the other hand, at the critical value of the Higgs mass we encounter a singularity of both anomalous scalings  $\mathbf{A}(t)$  and  $\mathbf{A}_I(t)$  at the instability point  $t_{\text{inst}}$  where  $\lambda(t_{\text{inst}}) = 0$ . This is depicted in Fig. 2 for  $\mathbf{A}_I(t)$  by the red curve corresponding to the slightly overcritical value 134.271 GeV (vertical red line and quasi-hyperbolic curves to the left and to the right of it). Other overcritical curves with small  $M_H > M_H^{\text{inst}}$  run through zero and change the sign of  $\mathbf{A}_I(t)$  from negative to positive. For larger  $M_H$ , when  $\lambda$  also becomes larger, the term  $-6\lambda$  in Eq. (3.9) for  $\mathbf{A}_I$  does not let it get positive at any  $t$ , and  $\mathbf{A}_I$  starts decreasing after reaching some maximal negative value (like for the green curve of  $M_H = 180$  GeV).

An important observation is that for all Higgs masses in the range  $M_H^{\text{inst}} = 134.27 \text{ GeV} < M_H < 185 \text{ GeV}$  the inflation range  $t_{\text{end}} < t < t_{\text{in}}$  is always below the instability value  $t_{\text{inst}} = 41.6$  (numerics gives that  $t_{\text{in}} < 34.8$ ), so that from Fig. 2  $\mathbf{A}_I(t)$  is always negative during inflation. Its running depicted in Fig. 2 explains the main difference from the results of the one-loop calculations in [3].  $\mathbf{A}_I(t)$  runs from big negative values  $\mathbf{A}_I(0) < -20$  at the electroweak scale to small but also negative values at the inflation scale below  $t_{\text{inst}}$ . This makes the CMB data compatible with the generally accepted Higgs mass range. Indeed,

the knowledge of the RG flow immediately allows one to obtain  $\mathbf{A}_I(t_{\text{end}})$  and  $x_{\text{end}}$  and thus find the parameters of the CMB power spectrum (3.16)–(3.18) as functions of  $M_H$ . The parameter of primary interest – the spectral index – is given by Eq. (3.16) with  $x = x_{\text{end}} \equiv N\mathbf{A}_I(t_{\text{end}})/48\pi^2$  and depicted in Fig. 3. Even for low values of the Higgs mass above the stability bound,  $n_s$  falls into the range admissible by the CMB constraint existing now at the  $2\sigma$  confidence level (based on the combined WMAP+BAO+SN data [26]),

$$0.94 < n_s(k_0) < 0.99. \quad (6.2)$$



**Figure 3:** The spectral index  $n_s$  as a function of the Higgs mass  $M_H$  for three values of the top quark mass.

The spectral index becomes too small (that is, dropping below 0.94) only for large  $x_{\text{end}}$  or large negative  $\mathbf{A}_I(t_{\text{end}})$ , which happens only when  $M_H$  either approaches the instability bound or exceeds 180 GeV at the decreasing branch of the  $n_s$  graph. Thus, we get lower and upper bounds on the Higgs mass, which both follow from a comparison of the *lower* bound of the spectral index in (6.2) with (3.16). Numerical analysis for the corresponding  $x_{\text{end}} \simeq -1.4$  gives for  $M_t = 171$  GeV the following range for a CMB-compatible Higgs mass:


$$135.62 \text{ GeV} \lesssim M_H \lesssim 184.49 \text{ GeV}. \quad (6.3)$$

Both bounds belong to the nonlinear domain of the equation (3.16) because their relevant  $x_{\text{end}} = -1.4 < -1$ . However, their calculation is still in the domain of our linear in logs approximation, because the quantity  $\mathbf{A}_I(t_{\text{end}})/16\pi^2 \simeq -0.07$  satisfies the restriction (5.12) for  $x_{\text{end}} = -1.4$ . For a smaller  $|\mathbf{A}_I(t_{\text{end}})|$  the expression (3.16) for  $n_s$  can be linearized in  $x$  and takes a particularly simple form,

$$n_s = 1 - \frac{2}{N} + \frac{\mathbf{A}_I(t_{\text{end}})}{48\pi^2}, \quad \mathbf{A}_I(t_{\text{end}}) \ll \frac{48\pi^2}{N} \sim 8. \quad (6.4)$$

It is applicable in a wide range of the Higgs mass in the interior of the domain (6.3).


As we see, the upper bound on  $n_s$  in (6.2) does not generate any restrictions on  $M_H$ , and it will not effect the  $M_H$ -range unless it will be lowered down by future CMB observations




**Figure 4:** The Einstein frame effective potential for the instability threshold  $M_H^{\text{inst}} = 134.27$  GeV. A false vacuum occurs at the instability scale  $t_{\text{inst}} \simeq 41.6$ ,  $\varphi \sim 80M_P$ , which is much higher than the inflation scale  $\varphi < \varphi_{\text{in}} \simeq 0.04M_P$ . A hypothetical inflation domain (ruled out by the lower  $n_s$  CMB bound and the requirement of the positive slope of  $\hat{V}$  at  $t \leq t_{\text{in}}$ ) is marked by dashed lines.

to about 0.964 — the top of the  $n_s$ -graphs in Fig. 3. The lower CMB bound in (6.3) is slightly higher than the instability bound  $M_H^{\text{inst}} = 134.27$  GeV. This bound depends on the initial data for weak and strong couplings and, what is even more important, on the top-quark mass  $M_t$  which is known with less precision. The bound  $M_H^{\text{inst}}$  given above was obtained for  $M_t = 171$  GeV. Below we consider basically this value of the top-quark mass. Results for the neighboring values  $M_t = 171 \pm 2$  GeV are presented in Fig. 3 only to show how strongly the plot gets shifted along the  $M_H$  axis. The general pattern of this shift follows the dependence of  $M_H^{\text{inst}}$  on  $M_t$  — the instability bound is larger for a larger top mass, which can be explained by a negative contribution  $-y_t^4 \sim -M_t^4$  to  $\mathbf{A}$ , and via  $\mathbf{A}$  to  $\beta_\lambda$ , (4.8).

Viability of the inflation scenario also implies that the system should safely evolve from the end of inflation to the EW vacuum through the periods of thermalization (reheating), radiation and matter domination. The main requirement for this is the possibility to roll down from  $\varphi_{\text{end}} = M_t \exp(t_{\text{end}})$  to  $\varphi = v$  or the positivity of the effective potential slope [4] (we disregard the tunneling scenario). The shape of the graph of this potential depicted in Fig. 4 for the instability threshold confirms the danger of having a negative slope for  $M_H$  close to  $M_H^{\text{inst}}$  — the formation of a false vacuum at the instability scale. At higher Higgs masses till about 160 GeV we first get a family of metastable vacua at scales  $\gtrsim t_{\text{inst}}$ . An example is the plot for the lower CMB bound  $M_H = 135.62$  GeV depicted in Fig. 5. For even larger  $M_H$  these metastable vacua get replaced by a negative slope of the potential which interminably decreases to zero at large  $t$  (at least within the perturbation theory range of the model), see Fig. 6. However, these metastable vacua and the negative slope of the potential are not dangerous for the inflationary scenario whose scales  $t_{\text{in}} < 34.8$  are



**Figure 5:** Inflaton potential at the lower CMB compatible value of  $M_H$ . A metastable vacuum exists at  $t \simeq 42$  which is much higher than the inflation domain at the positive slope of the potential.




**Figure 6:** The succession of effective potential graphs for  $M_H^{\text{inst}} < M_H < 184.3$  GeV showing the occurrence of a metastable vacuum followed for high  $M_H$  by the formation of a negative slope branch. Local peaks of  $\hat{V}$  situated at  $t = 34 \div 35$  grow with  $M_H$  for  $M_H \lesssim 160$  GeV and start decreasing for larger  $M_H$ .

much lower than the metastability scale and belong to the positive slope of  $\hat{V}$  to the left of its peak at  $t = 34 \div 35$ . Thus, even if the metastable vacuum occurs for low  $M_H$ , it exists before the inflation stage probed by current CMB observations<sup>12</sup>.

Let us finally focus on the running of  $\xi(t)$  depicted for five values of the Higgs mass

<sup>12</sup>The existence of this vacuum can perhaps be probed by wavelengths longer than that of a pivotal  $N \simeq 60$ , but this requires a deeper analysis.



**Figure 7:** Plots of running  $\xi(t)$ .

in Fig. 7, starting with the lower bound of the range (7.1) below. It is very slow for low values of the Higgs mass near the instability threshold, which of course follows from the smallness of the running  $\lambda(t)$  in this domain. Another property of the  $\xi$ -behavior is that the normalization of the power spectrum (5.8) leads to a value  $\xi \sim 5000$  for small Higgs masses, which is smaller than the old estimate  $\sim 10^4$  [7, 9, 1, 11, 21, 2, 3]. This is caused by a decrease of  $\lambda(t)$  which at  $t_{\text{in}}$  becomes much smaller than  $\lambda(0)$  – an observation made in [5]. This relaxes the situation with the “unnaturalness” problem in this model and brings  $\xi$  closer to a very subjective borderline between “natural” and “unnatural” values of coupling constants. For large  $M_H$  close to the upper bound of the CMB range,  $\xi(t)$  grows like  $\lambda(t)$  to the Landau pole and eventually goes out of the perturbation theory domain. However, it grows much faster than  $\lambda(t)$ , so that  $\hat{V}(\varphi) \sim \lambda(t)/\xi^2(t)$  tends to zero, as shown in Fig. 6 by the negative slope branches of the potential.

A final comment of this section concerns the change of numerical results for a “two-phase renormalizable” model of the system when the  $s$ -function is replaced by the step function, discussed in Sect. 4 as a justification for the modified RG scheme. This replacement leads to a negligible change in numerics – the instability threshold remains nearly the same, lower CMB bound on  $M_H$  grows by one in the second digit after the decimal point, while the upper bound decreases by 1 GeV.

## 7. Conclusions and discussion

We have found that the considered model looks remarkably consistent with CMB observations in the Higgs-mass range

$$135.6 \text{ GeV} \lesssim M_H \lesssim 184.5 \text{ GeV}, \quad (7.1)$$

which is very close to the widely accepted range dictated by electroweak vacuum stability and perturbation theory bounds.

Both bounds follow from the lower WMAP bound for the CMB spectral index  $n_s(k_0) > 0.94$  (the combined WMAP+BAO+SN data at the pivot point  $k_0 = 0.002 \text{ Mpc}^{-1}$  corresponding to  $N \simeq 60$  e-folds [26, 27]). The lower bound on  $M_H$  is very close to the instability threshold which in this model equals  $M_H^{\text{inst}} \simeq 134.27 \text{ GeV}$ . This is higher than the conventional SM estimate  $\sim 125 \text{ GeV}$ , which is caused by a strong non-minimal back reaction of gravity and/or the simplicity of our approximation disregarding higher loop orders.<sup>13</sup> Remarkably, the upper bound in (7.1) does not follow from a demand of a valid perturbation theory, but rather is imposed by the CMB data. This CMB mechanism works in a trustable perturbation regime with  $\lambda(t_{\text{in}}) < 2$ .

The current upper CMB bound  $n_s < 0.99$  does not impose restrictions on the Higgs mass, but it will generate bounds if the observed  $n_s$  essentially drops below the “classical” value  $n_s = 1 - 2/N \simeq 0.967$ . Conversely, this model will be falsified if the lower observational bound on  $n_s$  exceeds 0.964 — the top of the  $n_s$ -graphs in Fig. 3.

Our approach represents the RG improvement of our analytical results in [3]. In fact, here we completely recover the analytic formalism of [3] for all inflation parameters, which only gets modified by the RG mapping between the coupling constants at the EW scale and those at the scale of inflation. A peculiarity of this formalism is that for large  $\xi \gg 1$  the effect of the SM phenomenology on inflation is universally encoded in one quantity — the anomalous scaling  $\mathbf{A}_I$ . It was earlier suggested in [1] for a generic gauge theory (2.3), and in the SM it is dominated by the contributions of heavy particles —  $(W^\pm, Z)$ -bosons, top quark and Goldstone modes. This quantity is forced to run in view of RG resummation of leading logarithms, and this running raises a large negative EW value of  $\mathbf{A}_I$  to a *small negative* value at the inflation scale. Ultimately this leads to the admissible range of Higgs masses (7.1) very close to the conventional SM range.

In fact, this mechanism can be regarded as a kind of asymptotic freedom, for  $\mathbf{A}_I/64\pi^2$  determines the strength of quantum corrections in inflationary dynamics [11, 3]. Usually, asymptotic freedom is associated with the asymptotic decrease of  $\lambda(t)$  to zero. Here, this phenomenon is trickier because it occurs in the interior of the range (7.1) and fails near its lower and upper boundaries. For small Higgs masses in the range (7.1), as it follows from the plots of Fig. 1 and Fig. 2, running  $\lambda(t)$  gets very small when approaching the inflationary region  $t \sim 32 \div 35$  and generates a large  $\mathbf{A}_I(t)$  because of its  $1/\lambda$  part induced by vector boson and top quark particles. For large masses the coupling  $\lambda$  grows towards the Landau pole (though much slower than in nongravitating SM because of the suppression of the major part of the  $\lambda^2$ -term of its beta function). Therefore,  $\mathbf{A}_I(t)$  becomes big negative again, this time due to the negative contribution  $-6\lambda$  of the Goldstone modes, and falls out of the CMB range at  $M_H \simeq 185 \text{ GeV}$ .

Quantum effects are small only in the middle part of (7.1) with a moderately small  $\lambda$  where  $n_s$  is close to the “classical” limit  $1 - 2/N \simeq 0.967$  for  $x \equiv N\mathbf{A}/48\pi^2 \ll 1$ . Here

---

<sup>13</sup>It is well known that despite the perturbative range of coupling constants the two-loop RG improvement essentially lowers down the EW instability threshold compared to the one-loop RG running [43].

the original claim of [4] on smallness of quantum corrections is right, but this smallness, wherever it takes place, is achieved via a RG summation of big leading logarithms.

Qualitatively our main conclusions are close to those of [5] and [6], though the RG treatment in these works is very different from ours. In contrast to them, however, we did not try to suggest error bars for the Higgs mass range (7.1) and specify corrections due to the uncertainty in the top quark mass  $M_t$ . We only presented all our results for the middle value of the top mass  $M_t = 171$  GeV and demonstrated the sensitivity of the results to this choice by two plots of  $n_s$  with two neighboring values 169 GeV and 173 GeV (see Fig. 3). These error bars as well as the inclusion of the two-loop approximation are, in our opinion, not reliable. Generally speaking, all the conclusions of [5, 6] and our work still sound too good to be true in view of the conceptual problems that have not yet been resolved. These problems were partly mentioned in the criticism of [5] by Bezrukov and Shaposhnikov [6], and now we would like to briefly dwell on them.

As mentioned in [6], the RG flow of [5] depends (via the anomalous dimension of the Higgs field  $\gamma$ ) on the choice of gauge for local  $SU(2) \times U(1)$  invariance in the SM. This inalienable feature of any RG scheme in SM theory was bypassed in [6] as follows. The RG flow of all gauge-independent charges of the SM was used at the post-inflationary stage, while at the inflation stage the effect of running renormalization of the Higgs field (causing gauge dependence of the results) was disregarded due to the flatness of the effective potential  $\sim 1/\xi$ . In fact, this argument is misleading, because a small *gauge-dependent* slope of the inflation potential  $\sim \dot{\varphi}$  non-analytically enters the expression for the e-folding number  $N = \int d\varphi H(\varphi)/\dot{\varphi}$  – one of the main ingredients of the inflation parameters  $\zeta$  and  $n_s$ . Therefore, anyway, this brings an order  $O(1)$  gauge dependence into the expressions for these parameters.

In fact, this is a manifestation of a general problem of the off shell extension in quantum field theory. Unfortunately, we cannot formulate the calculation of CMB parameters as on-shell amplitudes or expectation values of physical observables uniquely defined at the quantum level. The power spectrum represents the quantum correlation function calculated at a special field-dependent moment of time – the horizon crossing. By using Heisenberg equations of motion (both for linear quantum perturbations and the nonlinear background) this function expresses as  $\zeta^2 = \hat{V}/24\pi^2 M_P^4 \hat{\varepsilon}$  entirely in terms of the inflaton potential  $\hat{V}(\varphi)$  and its gradients [32, 33, 34], cf. Eq.(3.10) for  $\hat{\varepsilon}$ . So the main inflation observable is the inflaton potential itself, but its quantum version  $\hat{V}^{\text{eff}}$  and its off-shell extension with  $(\hat{V}^{\text{eff}}(\varphi))' \neq 0$  – the effective potential – is gauge-dependent<sup>14</sup>. Moreover, off-shell extension of the effective potential (and more generally effective action) also depends on parametrization of quantum fields. A particular example of this dependence is given by different results for CMB characteristics obtained in different conformal frames – the original Jordan frame versus the Einstein one [3, 4, 6]. Another important example is the SM renormalization with different parameterizations of the Higgs multiplet. In the conventional quantization in Cartesian coordinates for the complex  $SU(2)$  doublet or real  $O(4)$  multiplet the theory is renormalizable and has one set of quantum corrections. In the

<sup>14</sup>In fact, this is a problem of consistent transition from  $\langle O[V(\varphi)] \rangle$  to  $O[V^{\text{eff}}(\langle \varphi \rangle)]$  which brings into game the problem of gauge dependence in the formalism of the mean field  $\langle \varphi \rangle$  and its effective action.

spherical coordinate system (used, for example, for the description of the chiral phase of SM [47, 6]) the theory is non-renormalizable and has another set of quantum corrections [48]. The S-matrices in both parameterizations coincide, but their off-shell effective actions are different.

Both off-shell aspects of the above type turn out to be important. The Einstein frame differs from the Jordan one by a field dependent conformal rescaling of the metric and a very nonlinear change of the Higgs field – radial variable of the Higgs multiplet. Quantization in these new variables has two major effects which are absent in the original Jordan frame. One effect is the change of the argument of the logarithms in quantum corrections, caused by conformal rescaling of mass parameters [3, 4, 22]. This renders these logarithms nearly constant and not contributing to quantum corrections. Another effect is the change of coefficients of these logs [35]. One can show that not only Higgs but also Goldstone contributions get suppressed in the pre-logarithm coefficients. Moreover, the Einstein frame used in [6] for the description of the chiral phase of the model automatically leads to the spherical parametrization of the Higgs multiplet, and this leads to an additional effect. The effective potential of the  $O(N)$  multiplet, calculated in spherical coordinates, does not contain Goldstone contributions even off shell<sup>15</sup>. Altogether, this leads to a big difference between the results in the Jordan and the Einstein frame – there are no large quantum corrections at high values of  $M_H$  when imposing the upper CMB bound on the Higgs range – a situation less sensitive to the CMB data, as claimed in [6].

This difference is interpreted in [6] as an inalienable theoretical ambiguity of the quantization scheme. However, we are not ready to completely endorse this statement and prefer a pragmatic approach which is to try to reduce this ambiguity as much as possible. Since the cosmological parameters  $\zeta$ ,  $n_s$ , etc. are the functionals of the effective potential, this attempt can be based on the choice of a preferred parametrization which automatically renders this potential gauge-independent. Such a parametrization indeed exists – this is spherical coordinates for the Higgs multiplet. These coordinates disentangle the only local gauge invariant observable, the radial variable  $\varphi = (\varphi^a \varphi^a)^{1/2} = (\Phi^\dagger \Phi)^{1/2}$ . The effective potential for this variable is gauge independent, so that this solves both parametrization<sup>16</sup> and gauge independence issues, but of course leaves open the problem of the conformal frame.

The price we will have to pay for such a resetting of the problem is that the theory becomes non-renormalizable even in the absence of gravitational interaction. Within such a setting it will make sense to include a two-loop RG improvement<sup>17</sup>, establish error bars

<sup>15</sup>On shell,  $V' = 0$ , the values of the effective potential of course coincide, because of a vanishing Goldstone mass (2.6).

<sup>16</sup>Of course, one might ask the question of effective potential dependence on the choice of parametrization for Goldstone angles, but we do not even want to tread into the discussion of this conundrum.

<sup>17</sup>Two-loop RG improvement and associated with it error bars in [6] most likely exceed available precision for a number of reasons. Not the least of them is that the two-loop potential was taken from the calculations of [49] in Cartesian coordinates, while its RG improvement in the chiral phase of inflating SM was based on beta functions derived in the spherical parametrization of the Higgs multiplet. Also, the conventional “Cartesian coordinates” RG flow of SM at low energy scales was matched with the chiral RG flow in spherical parametrization at the inflation stage. The consistency of this procedure requires checking both



associated with the current indeterminacy of the SM and CMB data and the indeterminacy in the post-inflationary cosmological scenario. All this goes beyond the goals of the present paper, and will be considered in future publications. Here we are forced to reconcile with the gauge dependence of the obtained results. Fortunately, this dependence which enters through the renormalization function  $Z_{\text{in}}$  in (5.8) is rather weak. The value of  $Z_{\text{in}}$  in the Feynman gauge differs from that of the Landau gauge (used throughout the paper) by about 6%, which ultimately effects the value of spectral index in the third digit after the decimal point.

Finally, in addition to a good match of the spectrum of cosmological perturbations with the CMB data our model also describes the mechanism of generating the cosmological *background* itself upon which these perturbations exist. This mechanism is based on the no-boundary or tunneling cosmological wavefunction [16, 17, 18, 19, 20] possibly prescribing initial conditions for inflation. Within the RG improvement the one-loop distribution function (2.4) [25, 1] for the no-boundary/tunneling state of the Universe is upgraded to  $\rho(\varphi) = \exp(\mp \Gamma(\varphi))$ , where  $\Gamma(\varphi)$  is the (on-shell) value of the Euclidean effective action of the quasi-de Sitter instanton, weighting in the quantum ensemble the member with the initial inflaton value  $\varphi$ . The calculation of this action gives  $\Gamma(\varphi) = -24\pi^2 M_P^4 / \hat{V}(\varphi)$  (see [11]) and the *tunneling* distribution function takes the form

$$e^{\Gamma(\varphi)} = \exp \left\{ -24\pi^2 \frac{M_P^4}{\hat{V}(\varphi)} \right\} = \exp \left\{ -96\pi^2 \frac{(M_P^2 + \xi(t)\phi^2)^2}{\lambda(t)\phi^4} \right\}. \quad (7.2)$$

For the CMB range of Higgs masses this distribution features sharp peaks at the scale  $\phi_0^2 \simeq \varphi_0^2 \simeq -64\pi^2 M_P^2 / \xi_0 \mathbf{A}_I(t_0)$ ,<sup>18</sup> see [50] for details. They exist only for  $\mathbf{A}_I(t_0) < 0$  – which is exactly our case – and, of course, correspond to the peaks of the potential in Fig. 6 at  $t = 34 \div 35$ . They have a very small quantum width  $\Delta\varphi/\varphi_0 \sim \sqrt{\lambda}/\xi$  and, therefore, can be interpreted as generating initial conditions for inflation at the quantum scale  $\varphi_0$  [1, 11, 21]. In the full CMB range of masses  $M_H$  their scale  $t_0$  exceeds the  $N = 60$  CMB formation scale  $t_{\text{in}}$ , as it should for sake of chronological succession of initial conditions for inflation and the formation of CMB spectra (in fact, this is a direct consequence of the positive slope of  $\hat{V}$  during inflation). Due to the high value of the exponentiated ratio  $\xi^2/\lambda$  in (7.2) the probability peak is very distinctive even for the lower CMB value of  $M_H = 135.6$  GeV. Moreover, for large  $M_H$  close to the upper CMB bound 185 GeV this peak gets separated from the non-perturbative domain of large over-Planckian scales due to a fast drop of  $\hat{V} \sim \lambda/\xi^2$  to zero shown on Fig. 6. This, in turn, follows from the fact that  $\xi(t)$  grows much faster than  $\lambda(t)$  when they both start approaching their Landau pole. Thus we get a complete *tunneling prescription* scenario of the quantum origin of the inflationary Universe [50].

To summarize, the inflation scenario driven by the SM Higgs boson with a strong non-minimal coupling to curvature looks very promising. This model supports the hypothesis

---

at the analytical and numerical levels.

<sup>18</sup>Naively this differs by sign from the one-loop scale of inflation (3.13), but in fact this is a nontrivial equation for  $\varphi_0$  because the right-hand side of this equality is a function of  $t_0$ .

that an appropriately extended Standard Model can be a consistent quantum field theory all the way up to the quantum-gravity scale and perhaps explain the fundamentals of all major phenomena in early and late cosmology [51, 52]. Ultimately, it will be the strongly anticipated discovery of the Higgs particle at LHC and a more precise determination of the primordial spectral index  $n_s$  by the Planck satellite that might decide the fate of this model.

## Acknowledgments

The authors are grateful to F. Bezrukov, M. Shaposhnikov and O. Teryaev for fruitful and thought-provoking correspondence and discussions and also benefitted from discussions with D. Diakonov, I. Ginzburg, N. Kaloper, I.M. Khalatnikov, D.V. Shirkov, S. Solodukhin, G.P. Vacca, G. Venturi and R. Woodard. A.B. and A.K. acknowledge support by the grant 436 RUS 17/3/07 of the German Science Foundation (DFG) for their visit to the University of Cologne. The work of A.B. was also supported by the RFBR grant 08-02-00725 and the grant LSS-1615.2008.2. A.K. and A.S. were partially supported by the RFBR grant 08-02-00923, the grant LSS-4899.2008.2 and by the Research Programme “Elementary Particles” of the Russian Academy of Sciences. The work of C.F.S. was supported by the Villigst Foundation. A.B. acknowledges the hospitality of LMPT at the University of Tours. A.S. also acknowledges RESCEU hospitality as a visiting professor.

## A. Renormalization of potential and non-minimal coupling terms – role of Higgs and Goldstone modes

Here we justify the  $s$ -factor suppression mechanism due to back reaction of non-minimally coupled gravitons. For simplicity we consider only the Higgs multiplet-graviton sector of the model and the renormalization of the potential and non-minimal coupling terms of the classical action

$$S = \int d^4x g^{1/2} \left( U(\Phi) R(g) - \frac{1}{2} (\nabla_\mu \Phi^a)^2 - V(|\Phi|) \right), \quad (\text{A.1})$$

$$U \equiv \frac{M_P^2 + \xi \Phi^2}{2}. \quad (\text{A.2})$$

For generality we work with the  $O(N)$  scalar multiplet  $\Phi^a$  which for  $N = 4$  corresponds to the real representation of the  $SU(2)$  complex Higgs doublet.

For renormalization of non-derivative terms of the action one expands it in quantum perturbations of the metric and scalar fields  $\sigma^A \equiv (h_{\mu\nu}, \sigma)$  on the curved background with a constant background scalar field  $\varphi^a$ ,  $\Phi^a = \varphi^a + \sigma^a$ ,  $\nabla_\mu \varphi^a = 0$ . To cancel nonminimal derivatives in the kinetic term of the action ( $\sim \nabla_\mu \nabla_\nu h^{\mu\nu}$ ) we pick up the background covariant DeWitt gauge-breaking term

$$S_{GB} = -\frac{1}{2} \int d^4x g^{1/2} U(\varphi) \left( \nabla^\mu h_{\mu\nu} - \frac{1}{2} \nabla_\nu h - \frac{\xi}{U(\varphi)} \varphi_a \nabla_\nu \sigma^a \right)^2. \quad (\text{A.3})$$

The quadratic part of the total action then takes the form

$$S_2 + S_{GB} = -\frac{1}{2} \int d^4x g^{1/2} \sigma^A \mathbf{F}_{AB} \sigma^B \quad (\text{A.4})$$

with the following operator acting in the space of gravitons  $h_{\mu\nu}$  and quantum Higgs multiplet  $\sigma^a$ , the index  $A$  running over the range  $A = (\mu\nu, a)$ ,

$$\mathbf{F}_{AB} = \begin{bmatrix} -UG^{\mu\nu, \gamma\delta} (\Box \delta_{\gamma\delta}^{\alpha\beta} + P_{\gamma\delta}^{\alpha\beta} + \frac{V}{U} \delta_{\gamma\delta}^{\alpha\beta}) & \frac{\xi}{2} g^{\mu\nu} \phi_b \Box + V_b^{\mu\nu} \\ \frac{\xi}{2} \phi_a g^{\alpha\beta} \Box + V_a^{\alpha\beta} & -\left(\delta_{ab} + \frac{\xi^2 \phi^2}{U} n_a n_b\right) \Box + \tilde{V}_{ab} \end{bmatrix}, \quad (\text{A.5})$$

where

$$P_{\mu\nu}^{\alpha\beta} = 2R_{(\mu}^{\alpha}{}_{\nu)}^{\beta} + 2\delta_{(\mu}^{(\alpha} R_{\nu)}^{\beta)} - \delta_{\mu\nu}^{\alpha\beta} R - g^{\alpha\beta} R_{\mu\nu} - R^{\alpha\beta} g_{\mu\nu} + \frac{1}{2} g^{\alpha\beta} g_{\mu\nu} R, \quad (\text{A.6})$$

$$V_a^{\mu\nu} = \frac{1}{2} g^{\mu\nu} n_a V' + (R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R) \xi \phi n_a, \quad (\text{A.7})$$

$$\tilde{V}_{ab} = m_G^2 (\delta_{ab} - n_a n_b) + m_H^2 n_a n_b - \xi \delta_{ab} R. \quad (\text{A.8})$$

Here, the term  $-\frac{\xi^2 \phi^2}{U} n_a n_b \Box$  in the kinetic part of  $\sigma^a$  originates from squaring the last term of the gauge, and  $m_G^2 = V'/\varphi$  and  $m_H^2 = V''$  are the Goldstone and Higgs masses given by Eqs. (2.6) and (2.7) above. They enter the potential term of the operator as coefficients of the projectors, respectively, transverse and longitudinal to the direction of the  $O(N)$ -multiplet vector of the background scalar field  $n^a = \varphi^a/\varphi$ ,  $\varphi \equiv \sqrt{\varphi^a \varphi^a}$ .

The matrix coefficient of  $\Box$  in  $\mathbf{F}_{AB}$  reads

$$\mathbf{C}_{AB} = \begin{bmatrix} -UG^{\mu\nu, \alpha\beta} & \frac{\xi}{2} g^{\mu\nu} \phi_b \\ \frac{\xi}{2} \phi_a g^{\alpha\beta} & -\delta_{ab} - \frac{\xi^2 \phi^2}{U} n_a n_b \end{bmatrix}, \quad (\text{A.9})$$

and its inverse equals

$$\mathbf{C}^{-1 AB} = \begin{bmatrix} -\frac{G_{\mu\nu, \alpha\beta}}{U} - \frac{1-s}{3U} g_{\mu\nu} g_{\alpha\beta} & \frac{s\xi\phi}{U} g_{\mu\nu} n_b \\ \frac{s\xi\phi}{U} n_a g_{\alpha\beta} & -(\delta_{ab} - n_a n_b) - s n_a n_b \end{bmatrix}, \quad (\text{A.10})$$

so that the operator  $\hat{\mathbf{F}} = \mathbf{C}^{-1} \mathbf{F}$  takes the form

$$\hat{\mathbf{F}} = \Box \hat{\mathbf{1}} + \hat{\mathbf{P}} - \frac{\hat{\mathbf{1}}}{6} R, \quad (\text{A.11})$$

where the potential term consists of the two pieces

$$\hat{\mathbf{P}} = \hat{\mathbf{P}}_\phi + \hat{\mathbf{P}}_R. \quad (\text{A.12})$$

The first term is independent of the curvature and all its elements are  $O(1/\xi)$  uniformly for the whole range of  $\varphi$  (or for all scales with  $1 \geq s > 1/6\xi$ ) except the element

$$(\hat{\mathbf{P}}_\phi)_b^a = -m_G^2 (\delta_b^a - n^a n_b) - m_H^2 s n^a n_b + n^a n_b O\left(\frac{1}{\xi}\right). \quad (\text{A.13})$$

It is dominated by the contribution of the Goldstone mass, and we explicitly present the term showing the suppression of the Higgs mass contribution by the factor  $s$ .

The second term is linear in the curvature and reads

$$\hat{\mathbf{P}}_R = \begin{bmatrix} P_{\mu\nu}^{\alpha\beta} + \frac{1}{6}R\delta_{\mu\nu}^{\alpha\beta} + O(\frac{1}{\xi}) & \frac{1-s-3s\xi^2\phi}{3U}Rg_{\mu\nu}n_b - \frac{2\xi\phi}{U}R_{\mu\nu}n_b \\ -s\xi\phi(R^{\alpha\beta} - \frac{1}{2}g^{\alpha\beta}R)n^a & (\hat{\mathbf{P}}_R)_b^a \end{bmatrix}, \quad (\text{A.14})$$

where

$$(\hat{\mathbf{P}}_R)_b^a = \left[ \left( \xi + \frac{1}{6} \right) (\delta_b^a - n^a n_b) + n^a n_b \left( s\xi - \frac{1}{6} + \frac{s}{3} \right) \right] R. \quad (\text{A.15})$$

Now we calculate the one-loop effective action

$$S^{1\text{-loop}} = \frac{i}{2} \text{Tr} \ln \left( \square \hat{\mathbf{1}} + \hat{\mathbf{P}} - \frac{\hat{\mathbf{1}}}{6} R \right) \quad (\text{A.16})$$

in the approximation linear in spacetime curvature.<sup>19</sup> Its divergent part is given by the trace of the second Schwinger-DeWitt coefficient  $\text{tr} \hat{\mathbf{a}}_2$  [53]. In this approximation it is just one half of the square of the full matrix potential term (A.12)<sup>20</sup>. Since the  $\hat{\mathbf{P}}_\phi$  part of this term commutes with the rest of the operator (A.11) in the  $1/\xi$ -approximation (in view of the matrix structure and in view of the assumed constancy of the background  $\nabla_\mu \varphi = 0$ ), it can be considered as a mass matrix

$$\hat{\mathbf{P}}_\phi = -\hat{\mathbf{M}}^2, \quad (\text{A.17})$$

so that

$$\hat{\mathbf{a}}_2 = \frac{1}{2} \hat{\mathbf{P}}^2 = \frac{1}{2} (\hat{\mathbf{M}}^2)^2 - \hat{\mathbf{P}}_R \hat{\mathbf{M}}^2 + \dots, \quad (\text{A.18})$$

and the one-loop action in dimensional regularization,  $d \rightarrow 4$ , takes the form explicitly featuring the Coleman–Weinberg structure of the effective potential,

$$\begin{aligned} S^{1\text{-loop}} = \int d^4x g^{1/2} & \left[ \frac{1}{32\pi^2} \left( \frac{1}{2 - \frac{d}{2}} - \mathbb{C} + \ln 4\pi \right) \text{tr} \hat{\mathbf{a}}_2 \right. \\ & - \frac{1}{64\pi^2} \text{tr} (\hat{\mathbf{M}}^2)^2 \left( \ln \frac{\hat{\mathbf{M}}^2}{\mu^2} - \frac{3}{2} \right) \\ & \left. + \frac{1}{32\pi^2} \text{tr} \hat{\mathbf{P}}_R \hat{\mathbf{M}}^2 \left( \ln \frac{\hat{\mathbf{M}}^2}{\mu^2} - 1 \right) + \dots \right]. \quad (\text{A.19}) \end{aligned}$$

<sup>19</sup>Because of the constant background scalar field, the Faddeev–Popov ghost contribution is  $\varphi$ -independent and does not contribute to the renormalization of the sector (A.1) of the model.

<sup>20</sup>With a non-constant background  $\nabla_\mu \varphi \neq 0$  the operator (A.11) acquires terms linear in derivatives, which can be absorbed into the generalized d'Alembertian by the redefinition of the covariant derivative,  $\nabla_\mu \rightarrow D_\mu = \nabla_\mu + O(\nabla_\mu \varphi)$ . Then, extra terms in  $\hat{\mathbf{a}}_2$  are given by the square of the curvature associated with the commutator of  $[D_\mu, D_\mu]$  and  $D_\mu^2 \hat{\mathbf{P}}$ . They are responsible for the renormalization of the kinetic term and curvature squared terms in the effective action, which we disregard here.

Here, indeed, the second line represents the Coleman-Weinberg potential and the third line gives the logarithmic renormalization of the non-minimal coupling, because after calculating the trace it takes the form  $U^{1-\text{loop}}(\varphi)R$ . Since  $\hat{\mathbf{M}}^2 \sim \varphi^2$ , the logarithms of the mass matrix yield, up to subleading corrections, a unit matrix  $\hat{\mathbf{1}} \ln \varphi^2$  and the traces equal

$$\begin{aligned} \frac{1}{64\pi^2} \text{tr} (\hat{\mathbf{M}}^2)^2 &= \frac{1}{32\pi^2} \left[ m_G^4(N-1) + s^2 m_H^4 \right] + O\left(\frac{1}{\xi}\right) \\ &= \frac{\lambda^2 \varphi^4}{128\pi^2} (6 + 18s^2) + O\left(\frac{1}{\xi}\right) \end{aligned} \quad (\text{A.20})$$

$$\begin{aligned} \frac{1}{32\pi^2} \text{tr} \hat{\mathbf{P}}_R \hat{\mathbf{M}}^2 &= \frac{1}{32\pi^2} R \left[ m_G^2(N-1) + s^2 m_H^2 \right] \xi + O(\xi^0) \\ &= \varphi^2 R \frac{3\lambda}{32\pi^2} (1 + s^2) \xi + O(\xi^0). \end{aligned} \quad (\text{A.21})$$

These equations show the suppression of the Higgs-mass contribution by the factors of  $s$  and the origin of  $6\lambda$  term of the anomalous scaling in Eq.(3.2) (cf. Eq.(2.8)). This term is entirely due to the contribution of Goldstone modes. Similarly, they explain the origin of the dominant part of the constant  $\mathbf{C} \sim \xi$  in (3.3), given by (3.5).

These equations also underlie the beta functions (4.8) and (4.9) in which we retained only the leading terms in  $\xi \gg 1$ , nontrivially depending on  $s$ -factors. One can check that subleading terms of (A.20) and (A.21) above are uniformly damped by inverse powers of  $\xi$  in the full range scales  $1 \geq s > 1/6\xi$ . They originate mainly due to graviton loops suppressed by powers of effective Planck mass  $1/(M_P^2 + \xi\varphi^2)$ . The remaining part of the beta functions (4.8)–(4.9) is generated from the heavy vector boson and quark sectors by adding to the mass matrix  $\hat{\mathbf{M}}^2$  their contribution  $\text{diag} [m_{W\pm}^2, m_Z^2, m_t^2]$  and by including the anomalous dimension terms caused by field renormalization.

## References

- [1] A. O. Barvinsky and A. Yu. Kamenshchik, Phys. Lett. B **332** (1994) 270.
- [2] F. L. Bezrukov and M. Shaposhnikov, Phys. Lett. B **659** (2008) 703.
- [3] A. O. Barvinsky, A. Yu. Kamenshchik and A. A. Starobinsky, JCAP **0811** (2008) 021.
- [4] F. L. Bezrukov, A. Magnon and M. Shaposhnikov, Phys. Lett. B **675** (2009) 88.
- [5] A. De Simone, M. P. Hertzberg and F. Wilczek, Phys. Lett. B **678** (2009) 1.
- [6] F. Bezrukov and M. Shaposhnikov, JHEP **0907** (2009) 089.
- [7] B. L. Spokoiny, Phys. Lett. B **147** (1984) 39.
- [8] T. Futamase and K.-I. Maeda, Phys. Rev. D **39** (1989) 399.
- [9] D. S. Salopek, J. R. Bond and J. M. Bardeen, Phys. Rev. D **40** (1989) 1753.
- [10] R. Fakir and W. G. Unruh, Phys. Rev. D **41** (1990) 1783.
- [11] A. O. Barvinsky and A. Yu. Kamenshchik, Nucl. Phys. B **532** (1998) 339.
- [12] E. Komatsu and T. Futamase, Phys. Rev. D **59** (1999) 0064029.
- [13] S. W. Hawking, Phys. Lett. B **115** (1982) 295.

- [14] A. A. Starobinsky, Phys. Lett. B **117** (1982) 175.
- [15] A. H. Guth and S.-Y. Pi, Phys. Rev. Lett. **49** (1982) 1110.
- [16] J. B. Hartle and S. W. Hawking, Phys. Rev. D **28** (1983) 2960; S.W. Hawking, Nucl. Phys. B **239** (1984) 257.
- [17] A. D. Linde, JETP **60** (1984) 211; Lett. Nuovo Cim. **39** (1984) 401.
- [18] V. A. Rubakov, JETP Lett. **39** (1984) 107.
- [19] Ya. B. Zeldovich and A. A. Starobinsky, Sov. Astron. Lett. **10** (1984) 135.
- [20] A. Vilenkin, Phys. Rev. D **30** (1984) 509.
- [21] A. O. Barvinsky and D. V. Nesterov, Nucl. Phys. B **608** (2001) 333.
- [22] M. Shaposhnikov and D. Zenhausern, Phys. Lett. B **671** (2009) 162.
- [23] J. Garcia-Bellido, D. G. Figueroa and J. Rubio, arXiv: 0812.4642 [hep-ph].
- [24] S. Weinberg, in *General Relativity*, ed. S. W. Hawking and W. Israel (Cambridge University Press, 1979) 790.
- [25] A. O. Barvinsky and A. Y. Kamenshchik, Class. Quant. Grav. **7** (1990) L181.
- [26] G. Hinshaw et al., Astrophys. J. Suppl. **180** (2009) 225.
- [27] E. Komatsu et al., Astrophys. J. Suppl. **180** (2009) 330.
- [28] S. Weinberg, *The quantum theory of fields. Vol.2. Modern applications*, CUP, Cambridge, 1996.
- [29] C. Amsler et al., “Particle Data Group”, Phys. Lett. **B667** (2008) 1
- [30] S. Coleman and E. Weinberg, Phys. Rev. D **7** (1973) 1888.
- [31] R. P. Woodard, Phys. Rev. Lett. **101** (2008) 081301, arXiv:0805.3089 [gr-qc].
- [32] V. F. Mukhanov and G. V. Chibisov, JETP Lett. **33** (1981) 532.
- [33] A. A. Starobinsky, Sov. Astron. Lett. **9** (1983) 302.
- [34] A. A. Starobinsky, Phys. Lett. B **91** (1980) 99.
- [35] A. O. Barvinsky, A. Yu. Kamenshchik and I. P. Karmazin, Phys. Rev. D **48** (1993) 3677.
- [36] J. L. F. Barbon and J. R. Espinosa, arXiv: 0903.0355 [hep-ph].
- [37] C. P. Burgess, H. M. Lee and M. Trott, arXiv: 0902.4465 [hep-ph].
- [38] F. Bezrukov, D. Gorbunov and M. Shaposhnikov, JCAP **0906** (2009) 029.
- [39] D. I. Kazakov, Theor. Math. Phys. **75** (1988) 440.
- [40] T.E. Clark, Boyang Liu, S.T. Love, T. ter Veldhuis, The Standard Model Higgs Boson-Inflaton and Dark Matter, arXiv:0906.5595v1 [hep-ph].
- [41] A. Sirlin and R. Zucchini, Nucl. Phys. B **266** (1986) 389.
- [42] R. Tarrach, Nucl. Phys. B **183** (1981) 384.
- [43] J. R. Espinosa, G. F. Giudice and A. Riotto, JCAP **0805** (2008) 002.
- [44] <http://www-theory.lbl.gov/~ianh/alpha.html>

- [45] D.V. Shirkov, Nucl Phys. **B 371**, 467 (1992); D. V. Shirkov and S. V. Mikhailov, Z. Phys. C **63** (1994) 463; R. S. Pasechnik, D. V. Shirkov and O. V. Teryaev, Phys. Rev. D **78** (2008) 071902.
- [46] M. Sher, Phys. Rept. **179** (1989) 273.
- [47] A.C. Longhitano, Phys. Rev. bf D22 (1980) 1166; S. Dutta, K. Hagiwara, Q.-S. Yan and K. Yoshida, Nucl. Phys. **B790** (2008) 111, arXiv:0705.2277.
- [48] P. F. Kelly, R. Kobes and G. Kunstatter, Phys. Rev. D **50** (1994) 7592 [arXiv:hep-ph/9406298]; R. Ferrari, arXiv:hep-th/0907.0426.
- [49] C. Ford, I.Jack and D. R. T. Jones, Nucl. Phys. **B387** (1992) 373; Erratum-ibid **B504** (1997) 551, hep-ph/0111190.
- [50] A. O. Barvinsky, A. Yu. Kamenshchik, C. Kiefer, A. A. Starobinsky, C. Steinwachs, work in progress.
- [51] T. Asaka, S. Blanchet and M. Shaposhnikov, Phys. Lett. B **631** (2005) 151; T. Asaka and M. Shaposhnikov, Phys. Lett. B **620** (2005) 17.
- [52] M. Shaposhnikov and D. Zenhausern, Phys. Lett. B **671** (2009) 187.
- [53] B. S. DeWitt, *Dynamical Theory of Groups and Fields*, Gordon and Breach, New York, 1965; A.O. Barvinsky and G.A. Vilkovisky, Phys. Reports, **119** (1985) 1.